

PEPSANE[®] ORAL GEL

GUAIAZULENE, DIMETHICONE

IN SINGLE-DOSE SACHETS

Read all of this leaflet carefully before taking this medicine. It contains important information about your treatment.

If you have any further questions or doubts, ask your doctor or pharmacist for additional information.

- Keep this leaflet, you may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If symptoms are getting worse or continue, please take medical advice.
- If you notice any side effects not listed in this leaflet or if you experience one of the side effects listed as serious, you must inform your doctor or your pharmacist.

In this leaflet:

1. What PEPSANE[®], oral gel in single-dose sachets is and what is it used for?
2. What do you need to know before taking PEPSANE[®], oral gel in single-dose sachets?
3. How to take PEPSANE[®], oral gel in single-dose sachets?
4. What are the possible side effects?
5. How to store PEPSANE[®], oral gel in single-dose sachets?
6. Other information

1. WHAT PEPSANE[®], oral gel in single-dose sachets IS AND WHAT IS IT USED FOR?

ANTIFLATULENT - ANTIACID

(A: Digestive system and metabolism)

This medicine is indicated for the treatment of stomach aches.

2. WHAT DO YOU NEED TO KNOW BEFORE TAKING PEPSANE[®], oral gel in single-dose sachets?

If your doctor told you that you have sugar intolerance, consult him/her before taking this medicine.

Do not take PEPSANE[®], oral gel in single-dose sachets:

- If you are allergic (hypersensitive) to one of the ingredients.

Some precautions must be taken with PEPSANE[®] oral gel in single-dose sachets:

Warning

This medicine contains parahydroxybenzoate and may cause allergic reactions.

IF YOU HAVE ANY DOUBTS, DO NOT HESITATE TO ASK YOUR DOCTOR OR YOUR PHARMACIST FOR ADVICE.

Taking other medicines

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including off-prescription medicines.

Pregnancy and breast-feeding

Under normal circumstances, this medicine may be taken during pregnancy or while breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

Excipients with known effects:

methyl parahydroxybenzoate, sorbitol.

3. HOW TO TAKE PEPSANE[®], oral gel in single-dose sachets?

Dosage

Oral route.

Take the contents of 1 or 2 single-dose sachets directly when the pain occurs, 2 to 3 times daily.

If you take more PEPSANE[®], oral gel in single-dose sachets than you should: Consult your doctor or pharmacist.

4. WHAT ARE THE POSSIBLE SIDE EFFECTS?

Like all medicines, PEPSANE[®], oral gel in single-dose sachets can cause side effects, although not everybody gets them.

Hypersensitivity reactions such as skin eruptions or pruritus have very rarely been reported.

If you experience any side effects not listed in this leaflet, or if the side effects become serious, inform your doctor or pharmacist.

5. HOW TO STORE PEPSANE[®], oral gel in single-dose sachets?

Keep out of the sight and reach of children.

Do not use PEPSANE[®], oral gel in single-dose sachets after the expiry date which is stated on the pack.

Store this medicine below 25°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. OTHER INFORMATION

What does PEPSANE[®], oral gel in single-dose sachets contain? The active substances are:

Dimethicone :3.000 g
Guaiazulene :0.004 g
In each sachet-dose

The other ingredients are:

Crystallising sorbitol 70%, carrageenates, sodium cyclamate, methyl parahydroxybenzoate, peppermint essential oil, purified water.

What is PEPSANE[®], oral gel in single-dose sachets and contents of the pack?

This medicine is available as an oral gel. Box of 12 or 30 single-dose sachets.

Marketing Authorisation holder

LABORATOIRES ROSA PHYTOPHARMA
68, RUE JEAN-JACQUES ROUSSEAU
75001 PARIS - FRANCE

This leaflet was last approved in January 2012.

Detailed information on this medicine is available on the ANSM (France) website.



Standard : +33 (0) 1 34 80 55 80
Medical information : + 33 (0) 1 34 80 72 60

7089920 - 403777